






BEDROCK GEOLOGY, SEARCH PHASE I PROJECT AREA, WESTERN SKEENA ARCH, WEST-CENTRAL BRITISH COLUMBIA




Detailed Geology of the Tatsi Creek Area




Detailed Geology of the Venture Peak Area




Detailed Geology of the Cariboo Mountain Area






Detailed Geology of the South Outcast Peak Area



Detailed Geology of the Limonite Creek Area



Detailed Geology of the Zymo Ridge Area



VOLCANIC AND SEDIMENTARY ROCKS

Lower to Upper Cretaceous

Skeena Group

Rocky Ridge Formation

LKRSFRV Dark green-grey to brown basalt to basaltic andesite flows and breccia; augite-, plagioclase- and hornblende-phyric to aphanitic, flow-banded rhyolite to dacite domes, breccia and lapilli tuff; intercalated sedimentary rocks including sandstone, siltstone and shale

Bulkley Canyon Formation

IKSBCS Sandstone, siltstone, mudstone and coal; sandstone is typically medium- to thick-bedded; local flaser bedding, laminated siltstone and sandstone; abundant plant fossils; ubiquitous detrital muscovite; local chert pebbles and volcanic-derived conglomerate and tuff

LKSV Undivided Skeena Group sedimentary and volcanic rocks

Jurassic to Cretaceous

Upper Jurassic to Lower Cretaceous

JKBS Undivided Bowser Lake Group and Skeena Group sedimentary rocks

Bowser Lake Group

Trout Creek assemblage

LJKBTCS Thick-bedded granule to boulder conglomerate, sandstone, calcareous sandstone, siltstone and shale; thick-bedded conglomerate with volcanic, plutonic and sedimentary rock derived clasts including sandstone and chert predominates; contains marine and nonmarine fossils

Netalzul volcanics

LJBNV Dark grey to green felspar- and hornblende-phyric andesite flows, breccia and tuff; intercalated mudstone, sandstone and pebble conglomerate; minor marine fossils

Muskaboo Creek assemblage

LJKMCS Sandstone, siltstone, and conglomerate; fine- to medium-grained, grey to brown to green sandstone in thin to thick planar beds predominates; interbedded with dark grey siltstone; rare granule to pebble conglomerate; common ripple marks, trough and hummocky cross-bedding, lenticular coquina and trace fossils

LKBS Undivided Bowser Lake Group sedimentary rocks

Triassic to Jurassic

Hazelton Group

Quock Formation

LJMHQS Thin-bedded dark grey siliceous mudstone and tan to pale pink tuff with felspar fragments; locally massive mudstone and lapilli tuff

Smithers Formation

LJMSH Tuffaceous sandstone, siltstone, shale and granule conglomerate; thin to thick planar beds of green to grey to brown sandstone and dark grey to black shale; abundant marine fossils

Saddle Hill Formation

LJMSHV Maroon to brick-red tuff; well-bedded crystal-lithic, ash and lapilli tuff and breccia; locally contains limestone fragments that weather recessively; minor aphanitic basalt

Niikitwa Formation

LJHNS Well-bedded tuffaceous shale, siltstone, greywacke and limestone

Upper Telkwa Formation

LJHT Flow-banded rhyolite and dacite domes, lapilli tuff, and crystal tuff; beige, cream, red, maroon, lavender and pink; spherulites locally up to 3 cm in diameter

LJHTD Red dacite tuff; well-bedded ash and lapilli tuff and breccia, rare coherent flows and welded tuff; minor rhyolite and aphanitic, vesicular basalt

LJHTB Black to dark green basalt flows and breccia; pyroxene- and plagioclase-phyric

LJHTA Andesitic breccia, lapilli-tuff, crystal tuff and flows; maroon to purple and plagioclase-phyric; minor basalt, dacite, rhyolite and volcanic derived sedimentary rocks

Lower Telkwa Formation

LJLHTB Coherent andesite flows; variably amygdaloidal and plagioclase-phyric, locally megacrystic; minor basalt and dacite

LJLHTA Andesitic breccia and lapilli tuff; monomict to polyimict; minor flows

LJLHTC Rhyolite flows and tuff; flow-banded coherent and lesser thin-bedded tuff

LJLHTD Polymict conglomerate, sandstone, siltstone and lapilli tuff

Undivided Telkwa Formation

LJLHTS Volcanic-derived sandstone, siltstone, minor conglomerate and tuff; well-bedded, green to maroon; includes well-bedded and massive reef limestone and calcareous sandstone and conglomerate

LJLHTA Undivided Telkwa Formation volcanic and sedimentary rocks

Upper Triassic

Stuhini Group

LJTS Argillite, siltstone and chert; black to dark grey and light grey to beige; typically thin-bedded

Mississippian to Permian

Zymoetz Group

Ambition Formation

LJZAS Limestone, marble, silty limestone, calcareous mudstone, minor green and pink lapilli tuff and volcanic sandstone; locally fossiliferous

Mount Attree volcanics

LJZASV Dark green andesitic to basaltic breccia, tuff and flows; aphyric to plagioclase- and augite-phyric; minor rhyolitic and dacitic breccia and tuff; intercalated with thin-bedded limestone, marble, calc-silicate, conglomerate, sandstone and mudstone

INTRUSIVE ROCKS

Eocene

Carpenter intrusive suite

ECl Granodiorite to granite; coarse-grained and equigranular; titanite, hornblende-, and biotite-bearing; minor rhyolite and andesite dyke complexes; undeformed to foliated

Nanika intrusive suite

ENGR Granite; fine-grained, plagioclase-, K-feldspar-, quartz-, biotite- and hornblende-porphyrific; beige to pale pink

ENd Diorite to gabbro; medium- to coarse-grained, equigranular; black and white; locally pegmatitic

ENi Undivided Nanika intrusive suite; granite, granodiorite and quartz monzonite; fine- to medium-grained, equigranular to hornblende-, biotite-, plagioclase- and/or quartz-porphyrific

Cretaceous to Eocene

KEI Undivided Cretaceous to Eocene intrusions; granodiorite, granite, quartz diorite, quartz monzonite, tonalite; fine- to medium-grained, equigranular to porphyritic

Late Cretaceous

Bulkley intrusive suite

LKbgd Granodiorite and quartz diorite; fine-grained, plagioclase- and hornblende-porphyrific; light to dark grey

LKbdi Diorite; fine-grained, equigranular to plagioclase-, hornblende- or pyroxene-porphyrific; typically biotite-bearing; dark green to grey

LKbqm Quartz monzonite, granite, quartz diorite and diorite; equigranular, medium- to coarse-grained; white, pink and black

LKbi Undivided Bulkley intrusive suite; granodiorite, quartz diorite, quartz monzonite, granite, and diorite; fine- to medium-grained, equigranular to plagioclase-, hornblende-, biotite-, quartz-, and/or K-feldspar-porphyrific

LKji Undivided Jurassic to Cretaceous intrusions in the CPC; diorite, gabbro and minor granodiorite; unfoliated to weakly foliated

Early Jurassic

Kleanza intrusive suite

EJkdj Diorite, quartz diorite, gabbro and monzonite; fine- to coarse-grained, equigranular to porphyritic; minor andesite dykes

EJkjd Granodiorite and granite; equigranular, medium- to coarse-grained

EJkpx Pyroxenite, equigranular, medium- to coarse-grained; locally layered

Undivided Kleanza intrusive suite

EJki Undivided Kleanza intrusive suite; granodiorite, granite, quartz monzonite, monzonite, quartz diorite, diorite, gabbro and pyroxenite; fine- to coarse-grained, equigranular to porphyritic

Late Triassic

Miligt intrusive suite

LTmqd Quartz diorite to diorite; fine- to coarse-grained, equigranular

LTmgd Granodiorite; fine- to medium-grained, plagioclase-, biotite- and hornblende-porphyrific

Mississippian to Permian?

Delta intrusive suite

MPDI Diorite, granodiorite, tonalite and gabbro; foliated to mylonitic

METAMORPHIC ROCKS

Paleozoic to Tertiary

Central Gneiss Complex

PZTCGM Amphibolite facies felsic and mafic volcanic rocks; tonalitic to granodioritic orthogneiss; minor marble and skarn; schistose and mylonitic

Geochronology (ages in Ma)

BC MINFILE Occurrences

- Subvolcanic Cu-Ag-Au (As-Sb)
- Intrusion-related Au pyrrhotite veins
- Epithermal Au-Ag; low sulphidation
- Porphyry Cu ± Mo ± Au
- Porphyry Mo (Low F-type)
- Cu skarn
- Au skarn
- Volcanic Redbed Cu
- Polymetallic veins Ag-Pb-Zn±Au
- Polymetallic manto Ag-Pb-Zn
- Noranda/Kuroko massive sulphide Cu-Pb-Zn
- Feldspar-quartz pegmatite

Geochronology (ages in Ma)

- K-Ar felspar
- K-Ar biotite
- K-Ar muscovite
- K-Ar hornblende
- U-Pb zircon
- U-Pb titanite

Structures

- Foliation or cleavage
- Fold hinge line
- Stretching lineation
- Bedding, tops unknown, right way up
- Bedding
- Igneous layering, flow banding

General

- River
- Road
- Contour line
- Lake
- Protected area

Faults / Contacts

- Thrust fault, defined / approximate / inferred / interpreted from geophysics
- Normal fault, defined / approximate / inferred / interpreted from geophysics
- Dextral fault, inferred / interpreted from geophysics
- Sinistral fault, interpreted from geophysics
- Unknown fault / strike-slip, defined / approximate / inferred / interpreted from geophysics
- Contact, defined / approximate / inferred / interpreted from geophysics
- Fold axial trace, inferred
- Outcrop Boundary

Magnetic declination

Year 2018
Latitude 54.5° N
Longitude 127.9° W
Model used IGRF12
18.12° E changing by 0.23° W per year

Geoscience BC
MAP 2019-03-01
MDRU MAP 17-2018
BCGS OPEN FILE 2019-07

BEDROCK GEOLOGY, SEARCH PHASE I PROJECT AREA, WESTERN SKEENA ARCH, WEST-CENTRAL BRITISH COLUMBIA

PART OF 1:250 000 NTS SHEETS 093L AND 103I

J.J. Angen, M. Rahimi, J.L. Nelson and C.J.R. Hart

1:150 000
UTM Zone 9; North American Datum 1983

February 2019

Product of the MDRU - Mineral Deposit Research Unit of The University of British Columbia with contributions by the British Columbia Geological Survey (BCGS) as part of Geoscience BC's SEARCH Project. This map comprises new mapping carried out during the summer of 2018 as well as interpretation of aeromagnetic data (Precision Geosurveys Inc., 2016) and compilation of previous mapping (Richards and Tipper, 1976; Handy, 1979; MacIntyre et al., 1989; Desjardins et al., 1990; Ryan, 1993; MacIntyre et al., 1994; Nelson et al., 2006; MacIntyre, 2007; Nelson et al., 2007; Everchick et al., 2008; Nelson et al., 2008; Nelson et al., 2008b; AMEC, 2009; Wasilenyevs and Yang, 2013). Geoscience BC is an independent, non-profit organization that generates earth science information in collaboration with First Nations, local communities, government, academia and the resource sector. Our independent earth science enables informed resource management decisions and attracts investment and jobs. Geoscience BC gratefully acknowledges the financial support of the Province of British Columbia, MDRU is a collaborative venture between the mining industry and The University of British Columbia. The unit, which operates on the support and financial assistance provided by the mining industry and the Natural Sciences and Engineering Research Council of Canada, is an internationally recognized research group devoted to solving mineral exploration-related problems. The BCGS is the government agency responsible for providing inventories, assessments, and archives of British Columbia's geology. It links government, the minerals industry, and communities to geoscience information and mineral resources. To stimulate investment and inform decisions about responsible land and resource management in the province, geoscience data are released through MapSpace, the BCGS online web service.

Recommended Citation: Angen, J.J., Rahimi, M., Nelson, J.L., and Hart, C.J.R. (2019). Bedrock geology, Search phase I project area, western Skeena arch, west-central British Columbia; Geoscience BC Map 2019-03-01, MDRU Map 17-2018 and BCGS Open File 2019-07, scale 1:150 000.